



**QUEEN'S
UNIVERSITY
BELFAST**

Living to the range limit: consumer isotopic variation increases with environmental stress

Reddin, C. J., O'Connor, N. E., & Harrod, C. (2016). Living to the range limit: consumer isotopic variation increases with environmental stress. *PeerJ*. <https://doi.org/10.7717/peerj.2034>

Published in:
PeerJ

Document Version:
Publisher's PDF, also known as Version of record

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

Copyright 2016 the authors.

This is an open access article published under a Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution and reproduction in any medium, provided the author and source are cited.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Living to the range limit: consumer isotopic variation increases with environmental stress

Carl J. Reddin¹, Nessa E. O'Connor² and Chris Harrod³

¹ Faculté des Sciences et des Techniques, Université de Nantes, Nantes, France

² School of Biological Sciences, Queen's University Belfast, Belfast, United Kingdom

³ Instituto de Ciencias Naturales Alexander Von Humboldt, Universidad de Antofagasta, Antofagasta, Chile

ABSTRACT

Background: Theoretically, each species' ecological niche is phylogenetically-determined and expressed spatially as the species' range. However, environmental stress gradients may directly or indirectly decrease individual performance, such that the precise process delimiting a species range may not be revealed simply by studying abundance patterns. In the intertidal habitat the vertical ranges of marine species may be constrained by their abilities to tolerate thermal and desiccation stress, which may act directly or indirectly, the latter by limiting the availability of preferred trophic resources. Therefore, we expected individuals at greater shore heights to show greater variation in diet alongside lower indices of physiological condition.

Methods: We sampled the grazing gastropod *Echinolittorina peruviana* from the desert coastline of northern Chile at three shore heights, across eighteen regionally-representative shores. Stable isotope values ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) were extracted from *E. peruviana* and its putative food resources to estimate Bayesian ellipse area, carbon and nitrogen ranges and diet. Individual physiological condition was tracked by muscle % C and % N.

Results: There was an increase in isotopic variation at high shore levels, where *E. peruviana*'s preferred resource, tide-deposited particulate organic matter (POM), appeared to decrease in dietary contribution, and was expected to be less abundant. Both muscle % C and % N of individuals decreased with height on the shore.

Discussion: Individuals at higher stress levels appear to be less discriminating in diet, likely because of abiotic forcing, which decreases both consumer mobility and the availability of a preferred resource. Abiotic stress might be expected to increase trophic variation in other selective dietary generalist species. Where this coincides with a lower physiological condition may be a direct factor in setting their range limit.

Subjects Ecology, Marine Biology, Zoology

Keywords Stress gradient, Intertidal, Range limit, Littorinid, Stable isotopes, Trophic niche width, Physiological condition, Diet, Trophic variation, Isotopic niche

INTRODUCTION

Environmental gradients, such as of temperature or moisture, are often associated with variation in a species' abundance and biomass (*Austin, 1985*), constraining its spatial

Submitted 12 March 2016

Accepted 22 April 2016

Published 1 June 2016

Corresponding author

Carl J. Reddin,
creddin01@qub.ac.uk

Academic editor

James Reimer

Additional Information and
Declarations can be found on
page 14

DOI 10.7717/peerj.2034

© Copyright

2016 Reddin et al.

Distributed under

Creative Commons CC-BY 4.0

OPEN ACCESS

range ([Normand et al., 2009](#); [Sexton et al., 2009](#)) and its potential contributions to ecosystem functioning ([Cardinale, Nelson & Palmer, 2000](#); [Duffy, 2002](#)). A species' spatial range can therefore expose the limits of its ecological niche ([Sexton et al., 2009](#)). However, most superficial indicators of species performance (e.g. abundance) do not help to identify whether a species' range limit is set directly or indirectly by the environmental variable(s) in question ([Somero, 2002](#); [Sexton et al., 2009](#)). Towards the extremes of a species' environmental tolerance (cf. Shelford's Law of Tolerance; [Odum & Barrett, 2005](#)), rates of biochemical reactions in individuals may be affected ([Menge & Sutherland, 1987](#)), and decreased individual performance may be observed as decreased physiological condition. An environmental gradient that is expected to produce a linear response in organism performance is the combined thermal and desiccation stress in intertidal ecosystems ([Lewis, 1964](#); [Somero, 2002](#)).

Typically a species' geographical range spans hundreds or thousands of kilometres, but vertical ranges may span only a few metres in the intertidal zone ([Connell, 1971](#); [Raffaelli & Hawkins, 2012](#)), where species are often distributed in conspicuous belts ([Stephenson & Stephenson, 1949](#); [Ingólfsson, 2005](#); cf. Continuum Concept, e.g. [Austin, 1985](#)). This tractable environmental gradient, identified as one of four major global ecoclines ([Whittaker, 1975](#)), forms as organisms that are principally marine compromise their physical and biological tolerances across height on the shore ([Connell, 1961](#); [Davenport & Davenport, 2005](#)), such that it can predict species richness and diversity (e.g. [Connell, 1978](#); [Zwerschke et al., 2013](#)). On the high shore, marine organisms are subjected to extreme abiotic stress from the synergistic effects of several physical factors combined ([Underwood, 1979](#); [Menge & Sutherland, 1987](#); e.g. desiccation, temperature; [Davenport & Davenport, 2005](#)), while other physical factors may oppose one another (e.g. wave exposure vs. desiccation; [Menge & Sutherland, 1987](#); [Harley & Helmuth, 2003](#); [Gilman, 2006](#)). The effect is that no single factor (e.g. elevation) drives the stress gradient, and integrated stress indicators, such as species dominance patterns, may be more useful in defining the gradient ([Menge & Sutherland, 1987](#)).

Components of a species' habitat can be quantitatively represented as geometric dimensions, brought together as potential niche space in a multi-dimensional hypervolume ([Hutchinson, 1978](#)). A species' theoretical optimum can be represented as a centroid or kernel in this niche space, with an individual suffering decreased performance when it is located far from this optimum ([Whittaker, Levin & Root, 1973](#); [Odum & Barrett, 2005](#)). A unimodal response in organism performance might then be expected around the optimum (fundamental niche). This [Hutchinson \(1978\)](#) concept of niche space is applied in stable isotope ecology, where isotopic axes represent n-space axes ([Bearhop et al., 2004](#); [Newsome et al., 2007](#)), with $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ typically representing variation in food web baseline (i.e. primary producers) and consumer trophic position, respectively. In particular, isotopic variance among individuals in the population can be informative regarding an organism's isotopic niche width, a close proxy for trophic niche width ([Bearhop et al., 2004](#); [Newsome et al., 2007](#); [Fink et al., 2012](#)). Innovative methods based on ecomorphological techniques allow the quantification and comparison of isotopic niche width ([Layman et al., 2007](#); [Jackson et al., 2011](#); [Syväranta et al., 2013](#)). Additionally,

the percentages of muscle C and N can provide proxies for physiological condition, with a decline in % C indicating a decline in stores of lipids and carbohydrates, and a decline in % N indicating a decline in tissue protein content (Nalepa et al., 1993).

Environmental stress gradients may be associated with changes in resource use; it might be expected that under ideal conditions a generalist but selective species may specialise on its preferred resource (Harrod, Mallela & Kahilainen, 2010), reducing niche width. However, environmental stress may decrease the abundance of the preferred resource or limit an organism's ability to forage (Hidalgo et al., 2008). This may encourage the assimilation of previously discounted resources (Harrod, Mallela & Kahilainen, 2010). Individuals may then be expected to either generalise their resource use similarly across the population (all feeding on equal proportions of different resources) or each individual to specialise their resource use variously (each feeding on distinct resources; Valen, 1965; Bearhop et al., 2004). Individual specialisation appears to be widespread among animal species (Bolnick et al., 2003; Araújo, Bolnick & Layman, 2011), and the degree of dietary specialisation of consumers has proven important in governing the strength of top-down control (Duffy et al., 2007). Still, few studies have calculated the magnitude of individual specialisation under different contexts (Araújo, Bolnick & Layman, 2011), such as a species' range limit.

We estimated the $\delta^{15}\text{N}$, $\delta^{13}\text{C}$, % N and % C of representative individuals of the regionally common grazing gastropod *Echinolittorina peruviana* (Lamarck, 1822) and its putative food sources. Individuals were sampled at three shore height levels at several rocky shores in northern Chile. Based on current theory (Hidalgo et al., 2008), we anticipated that individuals on the high shore, which approach the upper limit of this species distribution and a more harsh physical environment than the lower shore, would be forced to limit their foraging and only consume food resources that are immediately available. Littorinid snails are ubiquitous in the upper intertidal such that, historically, they were recommended for general definition of the supralittoral limit on shores worldwide (Stephenson & Stephenson, 1949; Lewis, 1964). Furthermore, littorinids are generalist but selective grazers, having a feeding preference for easily digested microalgae and ephemeral macroalgae (Lubchenco, 1978; Norton et al., 1990). We predicted that: (1) the isotopic niche width of *E. peruviana* will be broader in populations inhabiting the high shore compared to those lower on the shore (individuals forced into consuming distinct resources); and (2) the physiological condition indices of muscle % C and % N, representing proportion of carbon stores (lipids and carbohydrates) and protein content respectively (e.g. Nalepa et al., 1993), will be lower in high shore individuals relative to low shore individuals.

MATERIALS AND METHODS

Study region

The Humboldt Current system intertidal is well described (Santelices, Vasquez & Meneses, 1986; Fernández et al., 2000; Broitman et al., 2001), typically with dense beds of the mussel *Perumytilus purpuratus* on the mid shore (Broitman et al., 2001). Our study was conducted on the Mejillones Peninsula of northern Chile, close to the city of

Antofagasta (approx. 23°20'S, 70°30'W). On this coastline that forms the western extreme of the Atacama Desert, tidal emergence brings numerous disadvantages for intertidal organisms, including extremely high solar intensity (Antofagasta summer monthly average direct normal solar radiation = ~7000 Wh/m²; [EnergyPlus, 2013](#)), extremely low rainfall (Antofagasta mean = 7 mm year⁻¹; [Climatempo, 2013](#)) and warm air temperature (summer monthly mean = 20.1 ± 0.1 °C; [EnergyPlus, 2013](#)). The Mejillones Peninsula exemplifies strong variation in coastal upwelling, induced by coastal topography, and associated spatial variation in intertidal assemblage structure ([Marín, Delgado & Escribano, 2003](#); [Thiel et al., 2007](#); [Reddin et al., 2015](#)). [Reddin et al. \(2015\)](#) found *E. peruviana* abundance to have a strong negative association ($r = -0.74$, $P < 0.001$) with a proxy for the intertidal influence of upwelling. The upper shore is mainly dominated by *E. peruviana* (average density, 57 ± 99 individuals m⁻²), chthamoid barnacles and *Ulva* spp ([Nielsen & Navarrete, 2004](#)). The Antofagasta Bay intertidal is also inhabited by the invasive ascidian *Pyura praeputialis*, which was a dominant intertidal bioengineer, covering exposed rocks and increasing local invertebrate diversity ([Castilla et al., 2004](#)).

Field sampling

Because the absolute height and width of species' vertical ranges (i.e. that of *E. peruviana*) may vary corresponding to local conditions (e.g. [Stephenson & Stephenson, 1949](#)), we use the emergent biological feature of organism vertical zonation to define levels of integrated thermal and desiccation stress ([Menge & Sutherland, 1987](#)). Therefore, *E. peruviana* individuals were collected from the following zones: high shore shallow pools (< 10 cm) above all emergent macroalgae (high); above barnacle/*Perumytilus* zone (middle); and within or below barnacle/*Perumytilus* zone (low). *E. peruviana* individuals were not found above these zones. [Hidalgo et al. \(2008\)](#) found an average vertical distance of 0.35 m between these 'mid' and 'low' shore heights at the lower latitude of 11°46'S, in Peru.

E. peruviana individuals were sampled concurrently with [Reddin et al. \(2015\)](#), during February to March 2012, at the same sites ($n = 17$, Euclidean distance between the most northern and southern sites = ~83 km), which represented regional coastline variation (see [Reddin et al., 2015](#), for sites detail). Similar target sized individuals were collected (~12.5 mm, longest axis), aiming to minimise ontogenetic variation, and then pooled into low, mid and high groups ([Table 1](#)). Although there remained size variation among sampled individuals, this was not significantly different among the three shore height groups (one-way ANOVA for length, $MS = 0.04$, $F_{1,58} = 0.02$, $P = 0.89$). Shore height groups were balanced to the smallest group size ($n = 20$) by randomly dropping surplus individuals per height per site (i.e. maintaining regional representation).

We sampled the following putative trophic resources at all locations where *E. peruviana* were sampled: *Ulva* sp., brown macroalgae (either *Dictyota* sp. or *Lessonia nigrescens*; fresh growth only), epilithic biofilm (scraped from rocks using a spatula) and particulate organic matter (POM; a 5 l water sample filtered through a 0.7 µm Whatman GF/F filter until the filter showed a colour change). The only situation where littorinids had apparent access to all putative resources was on the low shore. At all heights, we predicted that POM residue from outgoing tides would remain available as a resource but at

Table 1 Mass and length (mean \pm SD) of *E. peruviana* individuals at three relative shore height groups.

Height	Mass (g)	Length (mm)	Samples (n)
Low	0.3 \pm 0.1	10.9 \pm 1.4	20
Mid	0.3 \pm 0.1	10.9 \pm 1.1	20
High	0.3 \pm 0.1	11.0 \pm 1.5	20

diminishing quantities with increased shore height; it would be doubtful that the other resources would occur in a thriving state. Biofilm would be present at a far thinner thickness on the mid-shore, and probably of a different flora, while macroalgal matter could drift in, including to the high shore. No other consistent and obvious resources were present at mid- and high shore heights.

We recorded individual littorinid wet-mass (in shell, to 0.1 g; mass range 0.1–0.7 g) and length (longest axis; to 0.1 mm), then dissected the muscle tissue (Lorrain *et al.*, 2002) and washed it with distilled water. Fresh macroalgal growth was isolated and rinsed with distilled water. POM samples were filtered through pre-combusted (550 °C for 5 h) 0.7- μ m Whatman GF/F filters. All samples were oven-dried at 65 °C for 48 h to a constant mass. Animal, biofilm and macroalgal samples were then homogenised and weighed (\pm 0.01 mg) into tin capsules (6 \times 4 mm, Sercon Ltd) on a Mettler Toledo XS3DU Microbalance, or cut into sections (GF/F filters). Biofilm samples were duplicated, with one duplicate prepared as above for $\delta^{15}\text{N}$ and the other decalcified to remove inorganic calcium carbonate contaminants (Ng, Wai & Williams, 2007). Decalcification proceeded by application of 10% hydrochloric acid drop by drop until bubbling ceased (Carabel *et al.*, 2006), after which samples were re-homogenised and standardised by dry mass into tin capsules as above.

Samples were combusted in a continuous-flow elemental analyser (CHNOS) interfaced to a mass spectrometer (IsoPrime 100, Center for Stable Isotope Biogeochemistry, University of California at Berkeley) to estimate $\delta^{13}\text{C}$, $\delta^{15}\text{N}$, and elemental % C and % N values. Isotope ratio data were expressed in the standard δ unit, as the ratio of heavy to light isotopes, in ‰ units: δ (‰) = $[(R_{\text{sample}}/R_{\text{reference}}) - 1] \times 10^3$, with $R = {}^{13}\text{C}/{}^{12}\text{C}$ for carbon and ${}^{15}\text{N}/{}^{14}\text{N}$ for nitrogen. Reference materials for the above calculation were the international standards of V-PDB for C and air for N. We used two calibration standards: the external standard ‘peach leaves,’ NIST SMR 1547, showed analytical precision to be 0.10 and 0.15 ‰ for $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$, and an internal standard, *Patella vulgata* muscle, suggested precision to be < 0.1 ‰ for both $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$.

Statistical analyses

We estimated isotopic niche width ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) by Bayesian ellipses (SEA.B), and also estimated carbon range (CR) and nitrogen range (NR) separately using a Bayesian implementation of the metrics by Layman *et al.* (2007; all SEAc, SEA.B, CR and NR from R package SIBER; Jackson *et al.*, 2011). CR and NR split the bivariate isotopic niche represented by the SEA.B into its univariate $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ axes. All estimates were

calculated by 10^4 posterior density draws and are reported using mode and 95% credible intervals. Statistical significance of differences between pairs of groups was estimated by comparing proportional overlap of posterior densities. We also tested for any interaction of vertical variation in isotopic niche width or physiology with oceanographic variation identified by [Reddin et al. \(2015\)](#), by geographically splitting individuals sampled from Antofagasta Bay (sites and Mejillones Peninsula (again, replicates balanced between geographical groups' shore height levels, by randomly dropping surplus replicates) and re-calculating SEA.B).

Monotonic relationships were tested for using Spearman's rank correlations, with 95% confidence intervals calculated using the R package 'psychometric' ([Fletcher, 2010](#)). We compared standard correlation significance values with those accounting for spatial autocorrelation via the [Dutilleul et al. \(1993\)](#), in R package 'Spatial Pack,' [Osorio & Vallejos, 2014](#)) method, which adjusts the degrees of freedom appropriately where spatial non-independence of data is detected. In no case did this change the level of significance so, for simplicity, we present the standard test results.

To estimate proportional contribution of the sampled putative resources to the diet of *E. peruviana*, we used the Bayesian mixing model SIAR ([Parnell et al., 2010](#)), with individuals split by shore height. We used the trophic fractionation values for the intertidal mollusc *Mytilus edulis* ([Dubois et al., 2007](#); $\delta^{13}\text{C} = 2.2 \pm 0.1$, $\delta^{15}\text{N} = 3.8 \pm 0.1$). To facilitate model runs, we performed a number of steps: (a) concentration dependence of resources was included; (b) macroalgal species, in particular *Ulva* sp., had overlapping $\delta^{13}\text{C}$ values, and so data were grouped a priori to model runs; (c) to represent geographical variation in the intertidal isotopic baseline, resources and consumers were split into Antofagasta Bay and Mejillones Peninsula following [Reddin et al. \(2015\)](#). Still, model runs frequently yielded large credible intervals for estimated resource contributions (particularly for the high shore) so we only report cases of significant differences in the same shore height group (within or between the Bay and Peninsula), as estimated by comparing proportional overlap of posterior densities. All analyses were implemented within the R statistical package ([R Development Core Team, 2008](#)).

RESULTS

Isotopic niche widths showed a marked positive trend with shore height ([Fig. 1](#)), with low shore individuals being isotopically most similar (SEA.B mode = 2.47 ‰^2 , 95% credible intervals = 1.61–3.88; significantly smaller than mid-shore SEA.B, probability of difference > 99.99%), mid-shore individuals having intermediate variation (mode 11 ‰^2 , 95% CI = 7.21–17.5; significantly smaller than high shore, probability > 95%) and high shore individuals being isotopically most dissimilar (SEA.B over eight times larger than that of low shore individuals; mode 21.6 ‰^2 , 95% CI = 13.9–33.4; significantly larger than low shore SEA.B, probability > 99.99%). This positive trend was shown for both C and N ranges ([Fig. 2](#)), with N range increasing most sharply, although credible intervals were large for these metrics (differences between shore heights were insignificant; probability < 95%). The positive trend between shore and SEA.B was observed over both Antofagasta Bay and Mejillones Peninsula, appearing more severe over the Mejillones




Figure 1 Isotopic variation among *Echinolittorina peruviana* individuals grouped by relative height on the shore. (A) Individual values on an $\delta^{15}\text{N}$ - $\delta^{13}\text{C}$ biplot, with standard ellipses (equivalent to a bivariate SD), corrected for small sample size (SEAc), plotted as calculated by the SIBER procedure (Jackson *et al.*, 2011). Plot lines and symbols represent shore height (see legend). (B) *E. peruviana* isotopic niche width (SEA.B) at different shore heights, plotted with 95, 75 and 50% credible intervals and mode as a black point. Macroalgae includes *Lessonia nigrescens*, *Dictyota* sp. and *Ulva* sp. pooled.




Figure 2 Separate (A) carbon and (B) nitrogen isotopic variation of *E. peruviana* individuals from different shore heights. Range estimates calculated by a Bayesian implementation of Layman *et al.* (2007) in the R package SIBER (Jackson *et al.*, 2011). A black dot shows the mode, while boxes represent the 95, 75 and 50% credible intervals.

Peninsula (Fig. 3) although no shore height group was significantly different in SEA.B between the bay and the peninsula. However, Mejillones Peninsula had significantly lower densities of *E. peruviana* (18 ± 57 individuals m^{-2}) relative to Antofagasta Bay (88 ± 115 individuals m^{-2} ; $P < 0.01$, t-test).




Figure 3 Comparison of isotopic niche widths (SEA.B) of *E. peruviana* individuals, grouped by relative height on shore, over Antofagasta Bay (black boxes) and Mejillones Peninsula (white boxes). This geographical split follows the ecological differences identified over this region by Reddin et al. (2015). Plotted are 95, 75 and 50% credible intervals (boxes) and the mode (black or white dot). Sample sizes per height level for both geographical groups, 'low' $n = 10$, 'mid' $n = 10$, 'high' $n = 7$.

We identified negative correlations between relative shore height and both individual muscle % C and % N (Fig. 4; % carbon $r_s = -0.34$, $df = 58$, $P < 0.01$, 95% confidence intervals = -0.55 – -0.10 ; % nitrogen $r_s = -0.52$, $df = 58$, $P < 0.0001$, 95% CI = -0.69 – -0.31), with a proportionally greater drop in % N than % C. This led to an increase in C:N ratio with increasing relative shore height (Fig. 4C; $r_s = 0.51$, $df = 58$, $P < 0.0001$; 95% CI = 0.31 – 0.69).

Mixing model derived dietary contribution estimates were most precise for low shore individuals (Fig. 5), having narrower credible intervals, while mid- and high shore contribution estimates overlapped considerably. On the low shore, POM (Bay mode = 0.59, 95% credible intervals = 0.41–0.73; Peninsula mode = 0.39, 95% CI = 0.22–0.53) contributed consistently more than biofilm (Bay mode = 0.25, 95% CI = 0.03–0.4; Peninsula mode = 0.15, 95% CI = 0.02–0.28; Fig. 5) in both bay and peninsula (both probability > 99%), while macroalgae (mode = 0.43, 95% CI = 0.24–0.73) contributed significantly more than biofilm (mode = 0.15, 95% CI = 0.02–0.28) on the peninsula (probability > 95%) and in the bay macroalgae contributed (mode = 0.1, 95% CI = 0–0.43) less than POM (mode = 0.59, 95% CI = 0.41–0.73; probability > 95%). On the mid shore, contribution estimates of biofilm were most precise, both Antofagasta Bay and Mejillones Peninsula having low modal values (Bay mode = 0.27, 95% CI = 0.01–0.46; Peninsula mode = 0.06, 95% CI = 0–0.33, lower than POM, probability = 94%), although differences between estimates for all resources were insignificant within mid- and high shore levels. Macroalgal contribution estimates were higher in peninsula locations than in the bay, for both low and mid-shores, but large credible intervals made this difference insignificant at the 5% level (probability = 94% between low shore estimates).




Figure 4 Trend of decreasing (A) % C and (B) % N, and (C) increasing C:N, of *E. peruviana* muscle tissue with increasing relative shore height. Shown are the median (dark line), inter-quartile range (box), range (whiskers) and outliers (points).

DISCUSSION

We found isotopic niche widths of an intertidal grazer to be significantly larger approaching the limit of its vertical distribution, coinciding with significant decreases in % C and % N, both candidate proxies for individual physiological condition (e.g. [Nalepa et al., 1993](#)). Vertical shore height is a well understood stress gradient for marine organisms ([Lewis, 1964](#); [Valdivia et al., 2011](#)) and governs their vertical distribution ([Stephenson & Stephenson, 1949](#); [Somero, 2002](#)). We document a case where individuals approaching their upper distribution limit, which is set by abiotic factors, may exhibit a more varied, suboptimal, potentially indiscriminating resource use, supported by a decrease in physiological condition and isotope mixing models. We compare evidence for this conclusion and for competing explanations of an enlarged isotopic niche width.

A large population isotopic niche width for this generalist species suggests that individuals within this population were isotopically distinct, and likely consumed isotopically distinct foods. Conversely, isotopically more similar individuals, showing lower isotopic variation at the population level, likely consumed isotopically similar (on average) foods ([Cummings et al., 2012](#)). The positive association between population isotopic niche width and putative abiotic stress therefore supports our hypothesis = that individuals at the high shore extreme of their environmental tolerance were forced to individually specialise their diet. *E. peruviana*, as well as other high shore littorinids, preferentially feeds on the epilithic microbial film ([Hawkins et al., 1989](#); [Norton et al., 1990](#); [Mak & Williams, 1999](#); [Hidalgo et al., 2008](#)), including bacteria, microalgae, macroalgal spores and early growth stages ([Norton et al., 1990](#); [Vermeulen et al., 2011](#)). *E. peruviana* can also consume crustose, foliose and filamentous macroalgae ([Santelices, Vasquez & Meneses, 1986](#)). We used the isotopic composition of muscle tissue, which has a turnover rate in marine molluscs of between 3–6 months ([Lorrain et al., 2002](#)).




Figure 5 Contributions of putative resources to *E. peruviana* diet at high (A and B), middle (C and D) and low (E and F) shore heights. Contributions estimated by SIAR mixing models (Parnell et al., 2010) run separately for Antofagasta Bay (A, C, E) and Mejillones Peninsula (B, D, F). Plotted are contribution estimates' 95, 75 and 50% Bayesian credible intervals (boxes), mode values (dots) and probability of differences between putative resource contributions of POM, macroalgae (*L. nigrescens*, *Dictyota* sp. and *Ulva* sp. pooled), and epilithic biofilm.

The isotopic composition of the assimilated resource is therefore averaged over that period, during which the mobile snail could move considerably, potentially even between the stress zones of this study, resembling other mobile species approaching their distributional limits. Benthic grazers can move considerably per tide to optimise their foraging, up to 1.5 m for the European littorinid, *Littorina littorea*, (Newell, 1958) and tidal vertical migrations of up to 1 m for a tropical limpet (Williams & Morritt, 1995). Therefore, we suggest that exchange of individuals between low and mid shore heights may be frequent but between mid and high levels would be low. Hidalgo et al. (2008) investigated the grazing of *E. peruviana* on wave-sheltered Peruvian shores, using two shore heights corresponding to 'low' and 'mid' heights in our study. They found that *E. peruviana* could control epilithic biofilm abundance better on the low shore and suggested that feeding was limited at greater height on the shore by faster desiccation of the rock's surface, decreasing the window for foraging per tidal cycle (also Mak & Williams, 1999; Norton et al., 1990). Additionally, greater quantities of mucus required to aid locomotion was suggested to burden the gastropod's energy reserves (Calow, 1974; Hidalgo et al., 2008). Littorinids have been reported to have higher rates of radular activity at higher shore locations, and higher shore species to feed more rapidly than low shore species (Newell, Pye & Ahsanullah, 1971). Foraging movement limitation may mean that littorinids in high shore habitats have to be more flexible in their diet, reflecting reduced resource availability. For instance, field observations of high-shore *E. peruviana* clustered over and leaving grazing scars on washed up kelp stipes support occasional consumption of macroalgae (see also Voltolina & Sacchi, 1990). In contrast, lower shore individuals may be able to move around and be more selective in their diet. Individual specialisation is widespread among animal species (Bolnick et al., 2003), including aquatic gastropods (Doi et al., 2010, using stable isotope analysis), although it has not been associated with approaching a species' spatial range limit.

We found both *E. peruviana* muscle % C and % N concentrations to decrease significantly with increasing putative stress. A relative decrease in % C is often interpreted as a decrease in tissue lipid content or carbohydrate storage (Nalepa et al., 1993), such as observed in freshwater molluscs during reproduction (Aldridge, 1982, via a decrease in C:N) or during low food availability (Russell-Hunter, Browne & Aldridge, 1984). Physiological processes resulting in the loss of muscle elementary N include high excretion of nitrogenous wastes, which increase with individual size, higher temperatures or longer durations, in the fish *Salmo trutta* (Elliott, 1976). Intertidal snails, however, need water to excrete their ammonia or urea. Protein catabolism was found to be higher in a high shore littorinid species relative to a low shore species (*Littorina saxatilis* and *Littorina obtusata*, respectively), but both had their highest rate of protein catabolism at lower temperatures (Aldridge, Russell-Hunter & McMahon, 1995). Protein catabolism and nitrogenous excretion could explain our observed % N decrease, but % N was not significantly related to $\delta^{15}\text{N}$ in our samples ($r_s = -0.17$, $P = 0.21$). Our individuals were size-matched, but both average temperature and the duration of emergence were expected to increase with height on the shore (Muñoz et al., 2008). High shore individuals were only found associated with small rock pools, presumably because of the shelter afforded

from thermal and desiccation stress. For instance, [Williams & Morritt \(1995\)](#) recorded individual limpet body temperatures from high shore shallow pools on Hong Kong Island to be consistently $\sim 2^\circ\text{C}$ cooler than exposed nearby limpets, although they noted that these pools could also become hypersaline or dry out after tidal emergence. Accordingly, [Hidalgo et al. \(2008\)](#) found that *E. peruviana* individuals inhabiting the high-shore were smaller, potentially due to less favourable conditions, although size-selective predation on the lower shore could also explain this. Whether our high shore individuals were there by intention or not is unclear but it is likely that the observed (by proxy) lower physiological condition might incur a lower fitness in high shore individuals, concordant with range limit conditions ([Sexton et al., 2009](#)).

We interpret our mixing model results to suggest that POM was the preferred food of *E. peruviana*, most probably that deposited on the rock during tidal immersion, which, we propose, becomes more scarce at greater shore height. The Humboldt region is characterised by high pelagic production ([Thiel et al., 2007](#)), which subsidises intertidal food webs ([Reddin et al., 2015](#)). *E. peruviana* was concluded by [Hidalgo et al. \(2008\)](#) to exert top-down control on biofilm as well as on crustose macroalgae, but had little effect on larger turf and canopy macroalgae. We suggest that our samples of biofilm, which were scraped to the underlying rock, may represent a resource that was too thick for the small littorinid's radula to fully penetrate, and that its grazing would have focussed on the surface layers of biofilm. A biofilm, especially with diatoms, can facilitate the deposition of suspended particles (i.e. POM) by production of an extracellular polymeric secretion (EPS; [Decho, 2000](#)). At higher shore heights, shorter tidal immersion time and greater UV radiation leads to less abundant biofilm ([Thompson, Norton & Hawkins, 2004](#)), itself leading to less deposited and captured POM. Together, combined with a shorter foraging time as the rocks dry during tidal emersion, could decrease *E. peruviana*'s consumption of preferred POM. Additionally, we observed an insignificant difference in rate of increase in isotopic variation with greater shore height, between Antofagasta Bay and Mejillones Peninsula. Higher POM concentrations in the waters of Antofagasta Bay ([Reddin et al., 2015](#)) may increase POM deposited with each tide, supporting high shore individuals. The higher abundance of ecosystem engineers in Antofagasta Bay (*P. purpuratus* and *Pyura praeputialis*; [Castilla et al., 2004](#); [Reddin et al., 2015](#); although see [Castilla et al., 2014](#)) could also ameliorate mid-intertidal conditions. Animals are expected to choose food resources that minimise energetic costs, including finding, handling and digesting the resource, thereby yielding the maximum 'value' per unit metabolic cost ([Townsend & Hughes, 1981](#)). This can be aided by behavioural maintenance of the animal's proximity to resources ([Underwood, 1979](#)).

The translation of isotopic niche to the trophic niche can be complicated by physiological conditions in the consumer or the resource(s). Stress caused by thermal extremes or water loss could itself contribute to the isotopic variation observed in high shore individuals. Intertidal organisms, including littorinids, have evolved behavioural and physiological tolerance mechanisms for greater desiccation and insolation ([Aldridge, Russell-Hunter & McMahon, 1995](#); [Somero, 2002](#); [Muñoz et al., 2008](#)). Some of these

physiological conditions can affect $\delta^{15}\text{N}$: excretion of nitrogenous waste in the form of urea, typical for higher intertidal littorinids (e.g. *Littorina saxatilis*; Aldridge, Russell-Hunter & McMahon, 1995), is associated with consumer enrichment in ^{15}N (Vanderklift & Ponsard, 2003); starvation or nutritional stress can result in elevated values of $\delta^{15}\text{N}$ (Vanderklift & Ponsard, 2003; Hobson, Alisauskas & Clark, 2013). A comparably high isotopic variation has nevertheless been observed previously in a grazing gastropod (the freshwater species *Lymnaea stagnalis*; Doi et al., 2010; $\delta^{13}\text{C}$ range 11.6 ‰, $\delta^{15}\text{N}$ range 7.4 ‰). Despite this variation being large relative to other sampled consumers, Doi et al. (2010) concluded that variation in resource use underpinned isotopic variation. Environmental temperature can affect trophic fractionation from resource to ectothermic consumer (Power, Guiguer & Barton, 2003; Barnes et al., 2007). Atmospheric temperature is not expected to vary across the sites but rock surface temperatures could vary depending on orientation to the sun, colour and type of rock. Water temperature was expected to differ by $\sim 3^\circ\text{C}$ between bay and outer peninsula situations (Reddin et al., 2015) but the resultant difference in trophic enrichment (+0.3 ‰ for both ^{13}C and ^{15}N , extrapolated from Barnes et al., 2007) was anticipated to be negligible (e.g. level of analytical error ± 0.1 ‰). We cannot rule out isotopic variation in the sources (Flaherty & Ben-David, 2010; Cummings et al., 2012); even if our consumer was feeding on the same resource(s) across shore heights a broader population isotopic niche width could result from variation in the basal resource(s). Thompson, Norton & Hawkins (2004) investigated biofilm biomass and diatom abundance at two different UK shore heights. They found strong negative correlations of biomass and diatom abundance with insolation stress and air temperature, but not with grazing intensity or dissolved nutrients (Thompson, Norton & Hawkins, 2004). Therefore, the fine-scale spatial variation of thermal and desiccation stress in the intertidal are likely to have a strong influence on biofilm biological and isotopic composition (Vermeulen et al., 2011), although this may be partially alleviated by the spatial averaging effect provided by mobile consumers (Post, 2002; Vermeulen et al., 2011). Cummings et al. (2012) recommend that the confounding effect of source variation is eliminated by transforming isotopic niche width to trophic niche width, by converting delta space to dietary proportions (p-space; Newsome et al., 2007). We were precluded from fully utilising this step, particularly on the high shore, because of difficulties identifying high shore resources and knowledge of the effect of environmental stresses at this level on isotopic variation.

When we combined our % C and % N values into the ratio C:N, we observed an increase in C:N ratio with increasing putative environmental stress via shore height. Increasing C:N is commonly used as a proxy for increasing organism condition, especially by lipid concentration (e.g. Schmidt et al., 2003), but such use has been criticised (Kidd et al., 2011). For example, beside condition, individual lipid content can vary with tissue type and stage in life history (e.g. reproductive stages). Furthermore, Kidd et al. (2011) found the relationship between C:N and lipid content to vary significantly among populations, with predictive models performing best within populations. An alternative interpretation of increasing C:N is a relative decrease in N, the variation of which can

exceed that of C (*Kidd et al., 2011*), as was the case in our study. Ultimately, a ratio is the relationship between two numbers; in the case of C:N, neither % C nor % N is solely dependent on the other, and an increase in the ratio can be derived from a change in either term. Where C:N is used as a proxy for physiological condition, we recommend that researchers support its appropriateness by presenting % N and % C, especially if processes that could change tissue % N are anticipated in their study system.

Environmental factors can induce limits on the effective feeding time of consumers, depending on their adaptations; for instance, ice or snow can impede grazing or predation in the high Arctic (*Connell, 1971*). Range limits, whether set by altitude in mountainous terrain, height on the intertidal or latitude across a continent, remain the spatial expression of the ecological niche (*Sexton et al., 2009*), suggesting that the upscaling of our results to other taxa, realms and stress scenarios may be possible. The optimal spatial scale of study depends on the acuteness of the stress gradient, but where exact replication is unlikely, variation in background processes should be integrated in the replication. This allows the averaging-out of the effects of alternative sympatric stress gradients (e.g. wave exposure in our study; see *Valdivia et al., 2011*) that may synergise or antagonise the effects of the focal stress gradient; thus, the conclusions develop generality. Therefore, the study extent, either spatial or temporal, must cover variation in regional (or seasonal) sources of stress, but not be so large that new food resources are introduced that could falsely inflate measures of trophic variation (*Araújo, Bolnick & Layman, 2011*). Conveniently, stable isotope ratios of organism tissues naturally provide a temporal (and spatial, depending on a species' mobility) averaging effect (*Post, 2002*), although the large foraging areas of highly mobile consumers may demand the adoption of large spatial grains.

ACKNOWLEDGEMENTS

We thank Joel Trexler for useful comments on an earlier version of the manuscript, two anonymous reviewers for comments that improved it further, and Felipe Docmac for help with sample collection.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

CH and CR were funded by the CONICYT Grant PAI MEL 81105006. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
CONICYT: PAI MEL 81105006.

Competing Interests

The authors declare that they have no competing interests.

Author Contributions

- Carl J. Reddin conceived and designed the experiments, performed the experiments, analyzed the data, wrote the paper, prepared figures and/or tables, reviewed drafts of the paper.
- Nessa E. O'Connor conceived and designed the experiments, wrote the paper, reviewed drafts of the paper.
- Chris Harrod conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, wrote the paper, reviewed drafts of the paper.

Data Deposition

The following information was supplied regarding data availability:

The raw data has been supplied as [Supplemental Dataset Files](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.2034#supplemental-information>.

REFERENCES

- Aldridge DW. 1982.** Reproductive tactics in relation to life-cycle bioenergetics in three natural populations of the freshwater snail, *Leptoxis carinata*. *Ecology* **63**(1):196–208 DOI [10.2307/1937044](https://doi.org/10.2307/1937044).
- Aldridge DW, Russell-Hunter WD, McMahon RF. 1995.** Effects of ambient temperature and of temperature acclimation on nitrogen excretion and differential catabolism of protein and nonprotein resources in the intertidal snails, *Littorina saxatilis* (Olivi) and *L. obtusata* (L.). *Hydrobiologia* **309**(1):101–109 DOI [10.1007/BF00014476](https://doi.org/10.1007/BF00014476).
- Araújo MS, Bolnick DI, Layman CA. 2011.** The ecological causes of individual specialisation. *Ecology Letters* **14**(9):948–958 DOI [10.1111/j.1461-0248.2011.01662.x](https://doi.org/10.1111/j.1461-0248.2011.01662.x).
- Austin MP. 1985.** Continuum concept, ordination methods, and niche theory. *Annual Review of Ecology and Systematics* **16**(1):39–61 DOI [10.1146/annurev.es.16.110185.000351](https://doi.org/10.1146/annurev.es.16.110185.000351).
- Barnes C, Sweeting CJ, Jennings S, Barry JT, Polunin NVC. 2007.** Effect of temperature and ration size on carbon and nitrogen stable isotope trophic fractionation. *Functional Ecology* **21**(2):356–362 DOI [10.1111/j.1365-2435.2006.01224.x](https://doi.org/10.1111/j.1365-2435.2006.01224.x).
- Bearhop S, Adams CE, Waldron S, Fuller RA, MacLeod H. 2004.** Determining trophic niche width: a novel approach using stable isotope analysis. *Journal of Animal Ecology* **73**(5):1007–1012 DOI [10.1111/j.0021-8790.2004.00861.x](https://doi.org/10.1111/j.0021-8790.2004.00861.x).
- Bolnick DI, Svanbäck R, Fordyce JA, Yang LH, Davis JM, Darrin Hulsey C, Forister ML. 2003.** The ecology of individuals: incidence and implications of individual specialization. *The American Naturalist* **161**(1):1–28 DOI [10.1086/343878](https://doi.org/10.1086/343878).
- Broitman BR, Navarrete SA, Smith F, Gaines SD. 2001.** Geographic variation of southeastern Pacific intertidal communities. *Marine Ecology Progress Series* **224**:21–34 DOI [10.3354/meps224021](https://doi.org/10.3354/meps224021).
- Calow P. 1974.** Some observations on locomotory strategies and their metabolic effects in two species of freshwater gastropods, *Ancylus fluviatilis* Mull and *Planorbis contortus* Linn. *Oecologia* **16**(2):149–161 DOI [10.1007/BF00345579](https://doi.org/10.1007/BF00345579).

- Carabel S, Godinez-Dominguez E, Verisimo P, Fernandez L, Freire J. 2006. An assessment of sample processing methods for stable isotope analyses of marine food webs. *Journal of Experimental Marine Biology and Ecology* 336(2):254–261 DOI 10.1016/j.jembe.2006.06.001.
- Cardinale BJ, Nelson K, Palmer MA. 2000. Linking species diversity to the functioning of ecosystems: on the importance of environmental context. *Oikos* 91(1):175–183 DOI 10.1034/j.1600-0706.2000.910117.x.
- Castilla JC, Guíñez R, Caro AU, Ortiz V. 2004. Invasion of a rocky intertidal shore by the tunicate *Pyura praeputialis* in the Bay of Antofagasta, Chile. *Proceedings of the National Academy of Sciences of the United States of America* 101(23):8517–8524 DOI 10.1073/pnas.0401921101.
- Castilla JC, Manríquez PH, Delgado A, Ortiz V, Jara ME, Varas M. 2014. Rocky intertidal zonation pattern in Antofagasta, Chile: invasive species and shellfish gathering. *PLoS ONE* 9(10):e110301 DOI 10.1371/journal.pone.0110301.
- Climatemps. 2013. Climate & Temperature. Available at <http://www.climatemps.com/> (accessed 20 July 2013).
- Connell JH. 1961. The influence of interspecific competition and other factors on the distribution of the barnacle *Chthamalus stellatus*. *Ecology* 42(4):710–723 DOI 10.2307/1933500.
- Connell J. 1971. On the role of natural enemies in preventing competitive exclusion in some marine animals and in rain forest trees. In: den Boer PJ, Gradwell GR, eds. *Proceedings of the Advanced Study Institute on "Dynamics of Numbers in Populations."* Oosterbeek: Centre for Agricultural Publishing and Documentation, 298–312.
- Connell JH. 1978. Diversity in tropical rain forests and coral reefs. *Science* 199(4335):1302–1310 DOI 10.1126/science.199.4335.1302.
- Cummings DO, Buhl J, Lee RW, Simpson SJ, Holmes SP. 2012. Estimating niche width using stable isotopes in the face of habitat variability: a modelling case study in the marine environment. *PLoS ONE* 7(8):e40539 DOI 10.1371/journal.pone.0040539.
- Davenport J, Davenport JL. 2005. Effects of shore height, wave exposure and geographical distance on thermal niche width of intertidal fauna. *Marine Ecology Progress Series* 292:41–50 DOI 10.3354/meps292041.
- Decho AW. 2000. Microbial biofilms in intertidal systems: an overview. *Continental Shelf Research* 20(10–11):1257–1273 DOI 10.1016/S0278-4343(00)00022-4.
- Doi H, Yurlova NI, Kikuchi E, Shikano S, Yadrenkina EN, Vodyanitskaya SN, Zuykova EI. 2010. Stable isotopes indicate individual level trophic diversity in the freshwater gastropod *Lymnaea stagnalis*. *Journal of Molluscan Studies* 76(4):384–388 DOI 10.1093/mollus/eyq020.
- Dubois S, Jean-Louis B, Bertrand B, Lefebvre S. 2007. Isotope trophic-step fractionation of suspension-feeding species: implications for food partitioning in coastal ecosystems. *Journal of Experimental Marine Biology and Ecology* 351(1–2):121–128 DOI 10.1016/j.jembe.2007.06.020.
- Duffy JE. 2002. Biodiversity and ecosystem function: the consumer connection. *Oikos* 99(2):201–219 DOI 10.1034/j.1600-0706.2002.990201.x.
- Duffy JE, Cardinale BJ, France KE, McIntyre PB, Thébault E, Loreau M. 2007. The functional role of biodiversity in ecosystems: incorporating trophic complexity. *Ecology Letters* 10(6):522–538 DOI 10.1111/j.1461-0248.2007.01037.x.
- Dutilleul P, Clifford P, Richardson S, Hemon D. 1993. Modifying the T test for assessing the correlation between two spatial processes. *Biometrics* 49(1):305–314 DOI 10.2307/2532625.
- EnergyPlus. 2013. Weather Data. Available at <https://energyplus.net/weather> (accessed 21 August 2013).

- Elliott JM. 1976. Energy losses in the waste products of brown trout (*Salmo trutta* L.). *Journal of Animal Ecology* 45(2):561–580 DOI 10.2307/3891.
- Fernández M, Jaramillo E, Marquet PA, Moreno CA, Navarrete SA, Patricio Ojeda F, Valdovinos CR, Vásquez JA. 2000. Diversity, dynamics and biogeography of Chilean benthic nearshore ecosystems: an overview and guidelines for conservation. *Revista Chilena de Historia Natural* 73(4):797–830 DOI 10.4067/S0716-078X2000000400021.
- Fink P, Reichwaldt ES, Harrod C, Rossberg AG. 2012. Determining trophic niche width: an experimental test of the stable isotope approach. *Oikos* 121(12):1985–1994 DOI 10.1111/j.1600-0706.2012.20185.x.
- Flaherty EA, Ben-David M. 2010. Overlap and partitioning of the ecological and isotopic niches. *Oikos* 119(9):1409–1416 DOI 10.1111/j.1600-0706.2010.18259.x.
- Fletcher TD. 2010. Psychometric: Applied Psychometric Theory. Available at <http://cran.r-project.org/package=psychometric>.
- Gilman SE. 2006. Life at the edge: an experimental study of a poleward range boundary. *Oecologia* 148(2):270–279 DOI 10.1007/s00442-006-0372-9.
- Harley CDG, Helmuth BST. 2003. Local- and regional-scale effects of wave exposure, thermal stress, and absolute versus effective shore level on patterns of intertidal zonation. *Limnology and Oceanography* 48(4):1498–1508 DOI 10.4319/lo.2003.48.4.1498.
- Harrod C, Mallela J, Kahilainen KK. 2010. Phenotype-environment correlations in a putative whitefish adaptive radiation. *Journal of Animal Ecology* 79(5):1057–1068 DOI 10.1111/j.1365-2656.2010.01702.x.
- Hawkins SJ, Watson DC, Hill AS, Harding SP, Kyriakidis MA, Hutchinson S, Norton TA. 1989. A comparison of feeding mechanisms in microphagous, herbivorous, intertidal prosobranchs in relation to resource partitioning. *Journal of Molluscan Studies* 55(2):151–165 DOI 10.1093/mollus/55.2.151.
- Hidalgo FJ, Firstater FN, Fanjul E, Cielo Bazterrica M, Lomovasky BJ, Tarazona J, Iribarne OO. 2008. Grazing effects of the periwinkle *Echinolittorina peruviana* at a central Peruvian high rocky intertidal. *Helgolander Marine Research* 62(S1):73–83 DOI 10.1007/s10152-007-0086-3.
- Hobson KA, Alisauskas RT, Clark RG. 2013. Stable-nitrogen isotope enrichment in avian tissues due to fasting and nutritional stress: implications for isotopic analyses of diet. *The Condor* 95(2):388–394 DOI 10.2307/1369361.
- Hutchinson GE. 1978. *An Introduction to Population Biology*. New Haven: Yale University Press.
- Ingólfsson A. 2005. Community structure and zonation patterns of rocky shores at high latitudes: an interocean comparison. *Journal of Biogeography* 32(1):169–182 DOI 10.1111/j.1365-2699.2004.01150.x.
- Jackson AL, Inger R, Parnell AC, Bearhop S. 2011. Comparing isotopic niche widths among and within communities: SIBER—Stable Isotope Bayesian Ellipses in R. *Journal of Animal Ecology* 80(3):595–602 DOI 10.1111/j.1365-2656.2011.01806.x.
- Kidd K, Fagan K-A, Koops MA, Arts MT, Power M. 2011. Assessing the utility of C:N ratios for predicting lipid content in fishes. *Canadian Journal of Fisheries and Aquatic Sciences* 68(2):374–385 DOI 10.1139/F10-119.
- Layman CA, Arrington DA, Montaña CG, Post DM. 2007. Can stable isotope ratios provide for community-wide measures of trophic structure? *Ecology* 88(1):42–48 DOI 10.1890/0012-9658(2007)88[42:CSIRPF]2.0.CO;2.
- Lewis J. 1964. *The Ecology of Rocky Shores*. London: EUP. 323 pp.

- Lorrain A, Paulet Y-M, Chauvaud L, Savoye N, Donval A, Saout C. 2002. Differential $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ signatures among scallop tissues: implications for ecology and physiology. *Journal of Experimental Marine Biology and Ecology* 275(1):47–61 DOI 10.1016/S0022-0981(02)00220-4.
- Lubchenco J. 1978. Plant species diversity in a marine intertidal community: importance of herbivore food preference and algal competitive abilities. *American Naturalist* 112(983):23–39 DOI 10.1086/283250.
- Mak YM, Williams GA. 1999. Littorinids control high intertidal biofilm abundance on tropical, Hong Kong rocky shores. *Journal of Experimental Marine Biology and Ecology* 233(1):81–94 DOI 10.1016/S0022-0981(98)00122-1.
- Marín V, Delgado LE, Escribano R. 2003. Upwelling shadows at Mejillones Bay (northern Chilean coast): a remote sensing in situ analysis. *Investigaciones Marinas* 31(2):47–55 DOI 10.4067/S0717-71782003000200005.
- Menge BA, Sutherland JP. 1987. Community regulation: variation in disturbance, competition, and predation in relation to environmental stress and recruitment. *American Naturalist* 130(5):730–757 DOI 10.1086/284741.
- Muñoz JLP, Camus PA, Labra FA, Finke GR, Bozinovic F. 2008. Thermal constraints on daily patterns of aggregation and density along an intertidal gradient in the periwinkle *Echinolittorina peruviana*. *Journal of Thermal Biology* 33(3):149–156 DOI 10.1016/j.jtherbio.2007.10.002.
- Nalepa TF, Cavaletto JF, Ford M, Gordon WM, Wimmer M. 1993. Seasonal and annual variation in weight and biochemical content of the zebra mussel, *Dreissena polymorpha*, in Lake St. Clair. *Journal of Great Lakes Research* 19(3):541–552 DOI 10.1016/S0380-1330(93)71240-X.
- Newell GE. 1958. The behaviour of *Littorina littorea* (L.) under natural conditions and its relation to position on the shore. *Journal of the Marine Biological Association of the United Kingdom* 37(1):229–239 DOI 10.1017/S0025315400014922.
- Newell RC, Pye VI, Ahsanullah M. 1971. Factors affecting the feeding rate of the winkle *Littorina littorea*. *Marine Biology* 9(2):138–144 DOI 10.1007/BF00348252.
- Newsome SD, Martínez del Río C, Bearhop S, Phillips DL. 2007. A niche for isotopic ecology. *Frontiers in Ecology and the Environment* 5(8):429–436 DOI 10.1890/060150.1.
- Ng JSS, Wai T-C, Williams GA. 2007. The effects of acidification on the stable isotope signatures of marine algae and molluscs. *Marine Chemistry* 103(1–2):97–102 DOI 10.1016/j.marchem.2006.09.001.
- Nielsen KJ, Navarrete SA. 2004. Mesoscale regulation comes from the bottom-up: intertidal interactions between consumers and upwelling. *Ecology Letters* 7(1):31–41 DOI 10.1046/j.1461-0248.2003.00542.x.
- Normand S, Treier UA, Randin C, Vittoz P, Guisan A, Svenning J-C. 2009. Importance of abiotic stress as a range-limit determinant for European plants: insights from species responses to climatic gradients. *Global Ecology and Biogeography* 18(4):437–449 DOI 10.1111/j.1466-8238.2009.00451.x.
- Norton TA, Hawkins SJ, Manley NL, Williams GA, Watson DC. 1990. Scraping a living: a review of littorinid grazing. *Hydrobiologia* 193(1):117–138 DOI 10.1007/BF00028071.
- Odum EP, Barrett GW. 2005. *Fundamentals of Ecology*. Belmont: Thomson Brooks/Cole.
- Osorio F, Vallejos R. 2014. SpatialPack: package for analysis of spatial data. R package version 0.2–3. Available at <http://cran.r-project.org/package=satialpack>.

- Parnell AC, Inger R, Bearhop S, Jackson AL. 2010. Source partitioning using stable isotopes: coping with too much variation. *PLoS ONE* 5(3):e9672 DOI 10.1371/journal.pone.0009672.
- Post DM. 2002. Using stable isotopes to estimate trophic position: models, methods, and assumptions. *Ecology* 83(3):703–718 DOI 10.2307/3071875.
- Power M, Guiguer KRRR, Barton DR. 2003. Effects of temperature on isotopic enrichment in *Daphnia magna*: implications for aquatic food-web studies. *Rapid Communications in Mass Spectrometry* 17(14):1619–1625 DOI 10.1002/rcm.1094.
- R Development Core Team. 2008. R: a language and environment for statistical computing. Vienne: R Foundation for Statistical Computing. Available at <http://www.R-project.org/>.
- Raffaelli D, Hawkins SJ. 2012. *Intertidal Ecology*. London: Springer Science & Business Media.
- Reddin CJ, Docmac F, O'Connor NE, Bothwell JH, Harrod C. 2015. Coastal upwelling drives intertidal assemblage structure and trophic ecology. *PLoS ONE* 10(7):e130789 DOI 10.1371/journal.pone.0130789.
- Russell-Hunter WD, Browne RA, Aldridge DW. 1984. Overwinter tissue degrowth in natural populations of freshwater pulmonate snails (*Helisoma trivolvis* and *Lymnaea palustris*). *Ecology* 65(1):223–229 DOI 10.2307/1939474.
- Santelices B, Vasquez J, Meneses I. 1986. Patrones de distribucion y dietas de un gremio de moluscos herbivoros en habitats intermareales expuestos de Chile central. *Monografías Biológicas* 4:147–171.
- Schmidt K, Atkinson A, Stubing D, McClelland JW, Montoya JP, Voss M. 2003. Trophic relationships among Southern Ocean copepods and krill: some uses and limitations of a stable isotope approach. *Limnology and Oceanography* 48(1):277–289 DOI 10.4319/lo.2003.48.1.0277.
- Sexton JP, McIntyre PJ, Angert AL, Rice KJ. 2009. Evolution and ecology of species range limits. *Annual Review of Ecology, Evolution, and Systematics* 40(1):415–436 DOI 10.1146/annurev.ecolsys.110308.120317.
- Somero GN. 2002. Thermal physiology and vertical zonation of intertidal animals: optima, limits, and costs of living. *Integrative and comparative biology* 42(4):780–789 DOI 10.1093/icb/42.4.780.
- Stephenson TA, Stephenson A. 1949. The universal features of zonation between tide-marks on rocky coasts. *Journal of Ecology* 37(2):289–305 DOI 10.2307/2256610.
- Syväranta J, Lensu A, Marjomäki TJ, Oksanen S, Jones RI. 2013. An empirical evaluation of the utility of convex hull and standard ellipse areas for assessing population niche widths from stable isotope data. *PLoS ONE* 8(2):e56094 DOI 10.1371/journal.pone.0056094.
- Thiel M, Macaya EC, Acuña E, Arntz WE, Bastias H, Brokordt K, Camus PA, Castilla JC, Castro LR, Cortés M, Dumont CP, Escibano R, Fernandez M, Gajardo JA, Gaymer CF, Gomez I, González AE, González HE, Haye PA, Illanes J-E, Iriarte JL, Lancellotti DA, Luna-Jorquera G, Luxoro C, Manriquez PH, Marín V, Muñoz P, Navarrete SA, Perez E, Poulin E, Sellanes J, Sepúlveda HH, Stotz W, Tala F, Thomas A, Vargas CA, Vasquez JA, Vega JMA. 2007. The Humboldt Current System of northern and central Chile: oceanographic processes, ecological interactions and socioeconomic feedback. *Oceanography and Marine Biology* 45:195–344.
- Thompson RC, Norton TA, Hawkins SJ. 2004. Physical stress and biological control regulate the balance between producers and consumers in marine intertidal biofilms. *Ecology* 85(5):1372–1382 DOI 10.1890/03-0279.

- Townsend CR, Hughes RN. 1981.** Maximising net energy returns from foraging. In: Townsend CR, Calow P, eds. *Physiological Ecology: An Evolutionary Approach to Resource Use*. Oxford: Blackwell Scientific Publishing, 86–108.
- Underwood AJ. 1979.** The ecology of intertidal gastropods. *Advances in Marine Biology* 16:111–210 DOI 10.1016/S0065-2881(08)60293-X.
- Valdivia N, Scrosati RA, Molis M, Knox AS. 2011.** Variation in community structure across vertical intertidal stress gradients: How does it compare with horizontal variation at different scales? *PLoS ONE* 6(8):e24062 DOI 10.1371/journal.pone.0024062.
- Valen L Van. 1965.** Morphological variation and width of ecological niche. *American Naturalist* 99(908):377–390 DOI 10.1086/282379.
- Vanderklift MA, Ponsard S. 2003.** Sources of variation in consumer-diet $\delta^{15}\text{N}$ enrichment: a meta-analysis. *Oecologia* 136(2):169–182 DOI 10.1007/s00442-003-1270-z.
- Vermeulen S, Sturaro N, Gobert S, Bouquegneau JM, Lepoint G. 2011.** Potential early indicators of anthropogenically derived nutrients: a multiscale stable isotope analysis. *Marine Ecology Progress Series* 422:9–22 DOI 10.3354/meps08919.
- Voltolina D, Sacchi CF. 1990.** Field observations on the feeding habits of *Littorina scutulata* Gould and *L. sitkana* Philippi (Gastropoda, Prosobranchia) of southern Vancouver Island (British Columbia, Canada). *Hydrobiologia* 193(1):147–154 DOI 10.1007/BF00028073.
- Whittaker RH. 1975.** *Communities and Ecosystems*. London: MacMillan.
- Whittaker RH, Levin SA, Root RB. 1973.** Niche, habitat, and ecotope. *American Naturalist* 107(955):321–338 DOI 10.1086/282837.
- Williams GA, Morritt D. 1995.** Habitat partitioning and thermal tolerance in a tropical limpet, *Cellana grata*. *Marine Ecology Progress Series* 124:89–103 DOI 10.3354/meps124089.
- Zwerschke N, Bollen M, Molis M, Scrosati RA. 2013.** An environmental stress model correctly predicts unimodal trends in overall species richness and diversity along intertidal elevation gradients. *Helgoland Marine Research* 67(4):663–674 DOI 10.1007/s10152-013-0352-5.